

TRIANUM-G

BIOLOGICAL FUNGICIDE



ACTIVE INGREDIENT:

Trichoderma harzianum Rifai strain KRL-AG2* 1.32%

OTHER INGREDIENTS

98.68%

TOTAL 100.00%

*Contains at least 1.0×10^7 colony forming units per gram dry weight.

KEEP AWAY FROM CHILDREN CAUTION

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, and then continue rinsing.
- Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information on (product, use, etc.), call the National Pesticides Information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific time (PT), seven days a week.
During other times, call the poison control center 1-800-222-1222.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation.

Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container under refrigerated conditions. Short periods at room temperatures above 75°F will not affect performance. Do not store near food or feed commodities. Do not store at temperatures above 75°F for prolonged periods. Keep container tightly closed when not in use.

DISPOSAL: Do not contaminate food or feed by disposal. Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. Do not reuse container. Dispose of empty container in a sanitary landfill, or by incineration, or by open burning if allowed by state and local authorities. If burned, keep out of smoke.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into the application equipment by shaking and tapping sides and bottom to loosen clinging particles, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration.

NET WEIGHT

44.1 lbs (20 kg)

EPA REG. NO.: 89635-2

EPA EST. NO.: 63119-NLD-001

LOT NUMBER: see packaging

EXPIRATION DATE: see packaging

MANUFACTURED BY

Koppert Biological Systems, Inc.

1502 Old US 23,

Howell, Michigan 48843 U.S.A.

PRODUCT NUMBER

12907



PERSONAL PROTECTIVE EQUIPMENT (PPE):

Mixer/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization. Follow manufacturer's directions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected workers may be in the area during application. For any requirement specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted - entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is protective eyewear, coveralls, waterproof gloves, shoes, and socks.

GENERAL INFORMATION

Triam-G is a preventative biological fungicide for control of soilborne plant pathogens and plant root diseases. The active ingredient is a microbe, *Trichoderma harzianum* Strain T-22, which when applied as an inoculant to soil, planting mixes or turf, grows onto plant roots as they develop and provides protection against plant root pathogens such as *Pythium*, *Rhizoctonia*, *Fusarium*, *Cylinrocladium* and *Thielaviopsis*.

Triam-G can be used in combination with or in rotation with other fungicides, insecticides, miticides, fertilizers or micronutrients in accordance with the more restrictive of label limitation and precautions. No label dosage rates should be exceeded. This product cannot be used with any product containing a label prohibition against such mixing. This product is not compatible with chemicals that contain the following active ingredients: benomyl, imazilil, propicanazole, febuconazole, and triflumizole. Do not apply Triam-G immediately before these pesticides are used. Where early season seed rot and seedling diseases are expected, use chemically treated seed or other appropriate measures for stand establishment and Triam-G for root disease control.

Note: Triam-G contains live spores of a microbe that must be used prior to disease onset. Strain T-22 becomes active in soil at temperatures above 50°F and is not effective while temperatures remain cold. Triam-G can be applied to sterilized or fumigated soil but must be applied after sterilization or fumigation.

This biological fungicide is for use in soil applications (drench, in soil furrow and potting soil) and seed treatments in or on all raw agricultural commodities, food and fiber crops, ornamentals, landscape plants, turf, and trees, including tree seedlings for transplanting into the forest.

USE RESTRICTIONS

Do not apply this product as a broadcast application. Cover applied product with soil or otherwise incorporate it into soil.

NOTE: DO NOT APPLY to sugarcane, pechay, rice, mushrooms, kiwi, tobacco, barley, oats, wheat, lemon, apple, and chickpea. Not for use on aquatic crops.

APPLY VIA GROUND APPLICATION ONLY.

CROPS	USE	RATE
Asparagus	In-furrow Greenhouse planting mix Nursery bed applications	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix 0.5 - 7.5 lbs. / 1000 sq. ft. incorporate with rake or tiller
Berries and Small Fruits: Blackberries, Blueberries, Currants, Elderberries, Gooseberries, Huckleberries, Loganberries, Raspberries, Strawberries, Grapes	In-furrow	2.5 - 6.0 lbs. / ½ acre
Bulb Crops: Garlic, Leeks, Onions, Shallots	In-furrow	2.5 - 6.0 lbs. / ½ acre
Citrus Fruits: Citrus Hybrids, Grapefruit, Kumquat, Limes, Oranges, Pummelos	In-furrow	2.5 - 6.0 lbs. / ½ acre
Cereal Grains: Buckwheat, Corn (grain, seed, sweet corn, silage, popcorn, high oil)	In-furrow	2.5 - 6.0 lbs. / ½ acre
Cucurbit Vegetables: Cucumbers, Melons, Gourds Pumpkins, Squash	In-furrow	2.5 - 6.0 lbs. / ½ acre
Flowers, Bedding Plants, and Ornamentals	In-furrow Greenhouse planting mix Nursery planting mix Nursery bed applications	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix 1.0 - 3.0 lbs. / cubic yd. soil mix 0.5 - 7.5 lbs. / 1000 sq. ft. incorporate with rake or tiller
Fruiting Vegetables: Eggplant, Sweet and Hot Peppers, Tomatillos, Tomatoes	In-furrow Greenhouse planting mix	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix
Herbs, Spices, and Mints	In-furrow Greenhouse planting mix	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix
Hydroponic Crops: Cucumbers, Tomatoes, Lettuce, Herbs and Spices	Greenhouse planting mix	1.0 - 3.0 lbs. / cubic yd. soil mix
Leafy Vegetables and Cole Crops: Arugula, Celery, Chervil, Endive, Fennel, Lettuce (head and leaf), Parsley, Radicchio, Rhubarb, Spinach, Swiss Chard, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Asparagus	In-furrow Greenhouse planting mix Nursery bed application	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix 1.0 - 3.0 lbs. / 1000 sq ft Incorporate with rake or tiller

CROPS	USE	RATE
Legume Vegetable (Succulent or Dried): Beans (soybeans, snap, dry), Lentils, Succulent and Dry Peas, Peanuts, Soybeans	In-furrow	2.5 - 6.0 lbs. / ½ acre
Oilseed Crops: Cotton, Canola, Safflower, Sunflower	In-furrow	2.5 - 6.0 lbs. / ½ acre
Pome Fruit: Pears, Quince	In-furrow Greenhouse planting mix Nursery planting mix Trees and shrubs	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix 1.0 - 3.0 lbs. / cubic yd. soil mix 0.2 oz. - 8.0 oz. / transplant hole and incorporate
Root Crops and Tuber Vegetables: Beets, Sugar beets, Carrots, Celeriac, Chicory, Horseradish, Parsnip, Radish, Rutabaga, Salsify, Turnips, Ornamental Roots and Tuber Vegetables	In-furrow	2.5 - 6.0 lbs. / ½ acre
Potatoes, Sweet Potatoes, Yams, Jerusalem Artichokes, Cassava, Ginger	Planting hole	0.2 - 8.0 oz. / transplant hole and incorporate
Shadehouse and Outdoor Nursery Crops: Deciduous Trees (Maple, Oak, etc.), Ornamentals, Grapes, Citrus, Pine	In-furrow Greenhouse planting mix Nursery planting mix Trees and shrubs	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix 1.0 - 3.0 lbs. / cubic yd. soil mix 0.2 - 8.0 oz. / transplant hole and incorporate
Stone Fruit: Apricots, Cherries, Nectarines, Peaches, Plums, Prunes	In-furrow Greenhouse planting Nursery planting mix Trees and shrubs	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix 1.0 - 3.0 lbs. / cubic yd. soil mix 0.2 - 8.0 oz. / transplant hole and incorporate
Tree Nuts: Almonds, Beech Nuts, Brazil Nuts, Butternuts, Cashews, Chestnuts, Filberts, Hickory Nuts, Macadamia Nuts, Pecans, Pistachios, Walnuts	In-furrow Greenhouse planting mix Nursery planting mix Trees	2.5 - 6.0 lbs. / ½ acre 1.0 - 3.0 lbs. / cubic yd. soil mix 1.0 - 3.0 lbs. / cubic yd. soil mix 0.2 - 8.0 oz. / transplant hole and incorporate
Turfgrass: Turf, sports fields, parks, established and new preparation of golf course tees, greens, fairways and roughs, seed production, sod production	New turf seedlings	Soil incorporate 1.0 - 4.0 lbs. / 1000 sq. ft. or 40 - 120 lbs./ acre

IN-FURROW SOIL TREATMENT: Apply Trianum-G from a granular applicator at the rate of 2.5 to 6 pounds per acre. For best results apply granules using a metering device, such as a Gandy applicator or equivalent, mounted to deliver granules into the furrow just behind the furrow opener. Apply in furrow at seeding or to established crops as a side dress.

NEW TURF SEEDINGS: Apply Trianum-G by ground application at 1 to 4 pounds per 1000 square feet or 40-120 pounds per acre just prior to or at the time of seeding. Incorporate into the soil surface or cover with mulch along with the grass seed. Application of irrigation water will help move Trianum-G into the soil.

GREENHOUSE PLANTING MIX AMENDMENT: Mix Trianum-G with planting mixes at a rate of 1 to 3 pounds per cubic yd. (loose) mix. For best results, thoroughly incorporate Trianum-G into planting mixes during mix preparation or at pot filling.

NURSERY PLANTING MIX AMENDMENT: For nursery planting mixes apply Trianum-G at the rate of 1 to 3 pounds per cubic yd. (loose) mix. For nursery beds, incorporate at a rate of 0.5 - 7.5 pounds per 1000 square feet of bed by raking or tilling. Application of irrigation water will help move Trianum-G into the soil. If soil is fumigated or steam sterilized apply Trianum-G after treatment.

TREES, SHRUBS, AND OTHER TRANSPLANTS: Apply Trianum-G granules to the planting hole at a rate of 0.2 to 8.0 ounces and incorporate into the planting soil or soil mix.

NOTICE TO BUYER AND SELLER

Seller warrants that this product conforms to the description on the label and is reasonably fit for the purposes stated on the label when used and stored in accordance with directions under normal conditions of use. To the extent permitted by state law, this warranty does not extend to use of this product contrary to label directions or under conditions not reasonably foreseeable by the Seller, and Buyer and User assume the risk of any such use. To the extent permitted by state law, Seller disclaims all other warranties express or implied, including any warranty of fitness or merchantability. To the extent permitted by state law, Seller shall not be liable for consequential, special or indirect damages resulting from use or handling of this product and Seller's sole liability and Buyer's and User's exclusive remedy shall be limited to refund of the purchase price. This product is sold only for uses stated on its label.

No express or implied license is granted to use or sell this product under any patent in any country except as specified.

EPA REG. NO. 89635-2

EPA EST. NO. 63119-NLD-001

MANUFACTURED BY

Koppert Biological Systems, Inc.

1502 Old US 23

Howell, Michigan 48843 U.S.A.





